


The Second Partial Test



Definition: Let (a, b) be a point contained in an open region R on which a function $z = f(x, y)$ is defined. Then (a, b) is a **critical point** if any of the following conditions are true:

1. $z_x(a, b) = 0 = z_y(a, b)$
2. $z_x(a, b)$ does not exist
3. $z_y(a, b)$ does not exist

Definition: Let (a,b) be a point contained in an open region R on which a function $z = f(x, y)$ is defined. Then (a,b) is a **critical point** if any of the following conditions are true:

1. $z_x(a,b) = 0 = z_y(a,b)$
2. $z_x(a,b)$ does not exist
3. $z_y(a,b)$ does not exist

Theorem: If $z = f(x, y)$ has a local maximum or a local minimum at a point (a,b) contained within an open region R on which $z = f(x, y)$ is defined, then (a,b) is a **critical point**.

Second Partial Test: Suppose $z = f(x, y)$ has continuous second partial derivatives on an open region containing a point (a, b) such that

$z_x(a, b) = 0 = z_y(a, b)$, and let

$$D = D(a, b) = \begin{vmatrix} z_{xx}(a, b) & z_{xy}(a, b) \\ z_{yx}(a, b) & z_{yy}(a, b) \end{vmatrix} = z_{xx}(a, b)z_{yy}(a, b) - z_{xy}(a, b)z_{yx}(a, b).$$

Second Partial Test: Suppose $z = f(x, y)$ has continuous second partial derivatives on an open region containing a point (a, b) such that


$z_x(a, b) = 0 = z_y(a, b)$, and let

$$D = D(a, b) = \begin{vmatrix} z_{xx}(a, b) & z_{xy}(a, b) \\ z_{yx}(a, b) & z_{yy}(a, b) \end{vmatrix} = z_{xx}(a, b)z_{yy}(a, b) - z_{xy}(a, b)z_{yx}(a, b).$$

Then:

1. If $D > 0$ and $z_{xx}(a, b) > 0$, $f(a, b)$ is a **local minimum**.
2. If $D > 0$ and $z_{xx}(a, b) < 0$, $f(a, b)$ is a **local maximum**.
3. If $D < 0$, $(a, b, f(a, b))$ is a **saddle point**.
4. If $D = 0$, the test is **inconclusive**.

Example: $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$



Example: $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

Example: $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$

Example: $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$

critical point



Example: $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$

critical point



$$z_{xx} = 4 \quad z_{xy} = 2$$

$$z_{yx} = 2 \quad z_{yy} = 2$$

Example: $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

critical point



$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$

$$z_{xx} = 4 \quad z_{xy} = 2$$

$$z_{yx} = 2 \quad z_{yy} = 2$$

$$D(-1,1) = \begin{vmatrix} 4 & 2 \\ 2 & 2 \end{vmatrix} = 8 - 4 = 4 > 0$$

$$z_{xx}(-1,1) = 4 > 0$$

Example: $z = f(x, y) = 2x^2 + 2xy + y^2 + 2x - 3$

$$z_x = 4x + 2y + 2$$

$$z_y = 2x + 2y$$

critical point




$$\begin{array}{l} 4x + 2y + 2 = 0 \\ 2x + 2y = 0 \end{array} \Rightarrow \begin{array}{l} 2x + y = -1 \\ x + y = 0 \end{array} \Rightarrow \begin{array}{l} x = -1 \\ y = 1 \end{array}$$

$$\begin{array}{ll} z_{xx} = 4 & z_{xy} = 2 \\ z_{yx} = 2 & z_{yy} = 2 \end{array} \quad D(-1,1) = \begin{vmatrix} 4 & 2 \\ 2 & 2 \end{vmatrix} = 8 - 4 = 4 > 0$$
$$z_{xx}(-1,1) = 4 > 0$$

Therefore, $f(-1,1) = -4$ is a **local minimum**,
and $(-1,1,-4)$ is a **minimum point**.

Try it now with $z = f(x, y) = x^3 - 3x + y^3 - 3y$!



Also try $z = f(x, y) = x^4 - y^4$ and $z = f(x, y) = x^4 + y^4$.

